

Research article

Published
2024-03-18

Cite as

Catherine E Taylor, Fidelis Masao, Jackson K Njau, Agustino Venance Songita and Leslea J Hlusko (2024) OH 89: A newly described ~1.8-million-year-old hominid clavicle from Olduvai Gorge, Peer Community Journal, 4: e34.

Correspondence

catherine_taylor@berkeley.edu
lesleahlusko@gmail.com

Peer-review

Peer reviewed and recommended by PCI Paleontology,

<https://doi.org/10.24072/pci.paleo.100218>



This article is licensed under the Creative Commons Attribution 4.0 License.

OH 89: A newly described ~1.8-million-year-old hominid clavicle from Olduvai Gorge

Catherine E Taylor^{ID,1}, Fidelis Masao^{ID,2,3}, Jackson K Njau^{ID,4,5}, Agustino Venance Songita³, and Leslea J Hlusko^{ID,1,6}

Volume 4 (2024), article e34

<https://doi.org/10.24072/pcjournal.372>

Abstract

Objectives Here, we describe the morphology and geologic context of OH 89, a ~1.8million-year-old partial hominid clavicle from Olduvai (Oldupai) Gorge, Tanzania. We compare the morphology and clavicular curvature of OH 89 to modern humans, extant apes, and a sample of other hominid fossil clavicles. **Materials and Methods** Comparative samples include 25 modern human clavicles, 30 *Gorilla*, 31 *Pan*, 7 *Papio*, and five hominid clavicles. Length regression on midshaft size using the extant comparative samples is used to estimate the total length of OH 89. A set of 9 linear measurements are taken from each individual. We also describe a new methodology for measuring clavicular curvature using measurements of sternal and acromial curvature, from which an overall curvature measurement is calculated. A principal component analysis (PCA) and a t-distributed stochastic neighbor embedding (tSNE) analysis are used to compare the morphology of OH 89 with the extant and fossil comparative samples. **Results** Our new method of measuring clavicular curvature successfully separates the different genera of the extant clavicles. The length estimate and sternal and acromial curve measurements for OH 89 falls within the larger male humans. The PCA shows OH 89 and most of the fossil hominids falling between the modern human and *Pan* groups, while the t-SNE suggests that OH 89, KSD-VP-1/1, KNM-ER 1808, and OH 48 are more similar to each other than to any of the other groups. This analysis also plots KNM-WT 15000 with the modern humans and Krapina 158 with the *Pan* individuals. **Discussion** The OH 89 clavicle derives from an individual of unknown hominid species with a shoulder breadth similar to that of a large human male. The curvature of OH 89 is relatively human-like relative to its length. Our new methodology for measuring clavicular curvature, combined with the utilization of t-SNE analyses and comparison of t-SNE results to PCA results, provides greater separation of genera than previously used methods, and wider use of t-SNE may be useful in paleoanthropological work.

¹Human Evolution Research Center, University of California Berkeley – Berkeley, CA, USA, ²University of Dar es Salaam – Dar es Salaam, Tanzania, ³Conservation Olduvai Project – Dar es Salaam, Tanzania, ⁴Department of Earth and Atmospheric Sciences, Indiana University – Bloomington, Indiana, USA, ⁵The Stone Age Institute – Bloomington, Indiana, USA, ⁶Centro Nacional de Investigación sobre la Evolución Humana – Burgos, Spain



Introduction

Until recently, paleoanthropologists did not dedicate much attention to the evolution of the hominid clavicle because its morphology is highly variable, and clavicles are relatively poorly preserved in the fossil record compared to other long bones. There are only a handful of decently preserved clavicles from Miocene apes, including a nearly complete *Nacholapithecus* clavicle from Kenya (Senut et al., 2004). The oldest hominid clavicle known to date is a fragment from *Ardipithecus* from the Afar region of Ethiopia (5.8-5.2Ma; Lovejoy, Suwa, Simpson, Matternes, & White, 2009). The Afar of Ethiopia has also yielded a half dozen clavicular remains of *Australopithecus afarensis*, including a left clavicle associated with the partial skeleton KSD-VP-1/1 (3.6 Ma; Haile-Selassie et al., 2010; Melillo, 2016), both clavicles of a juvenile partial skeleton DIK-1-1 (Alemseged et al., 2006), and a handful of fragmentary clavicles from Hadar dating to 3.0-3.2Ma (Drapeau, Ward, Kimbel, Johanson, & Rak, 2005; Lovejoy, Johanson, & Coppens, 1982). *Australopithecus* clavicles are also known from South Africa. Sterkfontein has yielded partial clavicles StW 431g, StW 582, and StW 606, as well as the StW 573 specimen that preserves both clavicles of an *Australopithecus* individual dated to ca. 3.67 Ma (Carlson et al., 2021; Green, 2020). Two specimens of *Australopithecus sediba* (MH1 and MH2) from the site of Malapa also preserve incomplete clavicular fragments (1.95 Ma; Dirks et al., 2010). Although there are in total just over a dozen known clavicles of representing *Ardipithecus* and *Australopithecus*, these are mostly fragmentary and consequently, difficult to study.

As for the genus *Homo*, as we approach *H. sapiens* in time, the number of fossil clavicles increases. For example, KNM-WT-15000 from northern Kenya preserves nearly complete right and left clavicles of a juvenile *Homo erectus* (~1.6 Ma; Walker, Leakey, & Leakey, 1993). KNM-WT 15000 has a notably short clavicle for its overall body size, though not outside of the range of modern human variation (Roach & Richmond, 2015). There are multiple partial clavicles attributed to *Homo naledi*, including U.W. 101-258, U.W. 101-1229, and U.W. 101-1347 (Feuerriegel et al., 2017). The majority of fossil clavicles are found in Europe, from Krapina, La Ferrassie, El Sidron, Gran Dolina, Sima de los Huesos, Kebara, Régardou, La Chapelle aux-Saints (Carretero, Arsuaga, & Lorenzo, 1997; Ferna et al., 1999; García-González, Rodríguez, Salazar-Fernández, Arsuaga, & Carretero, 2023; Gomez-Olivencia et al., 2018; Heim, 1982; Radović & Wolpoff, 1988; Rosas et al., 2016; Trinkaus, 2011; Voisin, 2006). Sima de las Palomas has also yielded a nearly complete Neanderthal clavicle (Walker et al., 2011). There are also roughly two dozen clavicles from Taforalt (Morocco), one from Qafzeh, and one from Shanidar (Trinkaus, 1982; Vandermeersch, 1981; Voisin 2006). There are multiple clavicles from Dmanisi, including the nearly complete D2724 (Lordkipanidze et al., 2007). For the earliest representatives of *Homo sapiens*, there is the Omo 1 partial skeleton that preserves a complete left clavicle from ~130 ka (Day, 1969; Pearson, Royer, Grine, & Fleagle, 2008; Voisin, 2008).

The discovery of the clavicle from partial skeleton of *Australopithecus afarensis* KSD-VP-1/1 brought renewed attention to the clavicle and interpreting its role in the evolution of the hominid shoulder girdle (e.g., Melillo, 2016; Melillo et al., 2019). Applying new methodological approaches, Melillo (2016) concluded that all *Australopithecus* clavicular remains are indistinguishable from each other and that the clavicle of *A. afarensis* has morphological affinities to both chimpanzees and humans.

Although clavicles themselves do not provide as clear a taxonomic or functional signal as other aspects of the skeleton, the relatively conserved nature of the primate shoulder girdle does provide information about the evolution of primate locomotion in a broader sense. For example, from a much wider taxonomic perspective, the bones that make up the stylopod, zeugopod, and autopod of tetrapod forelimbs are fairly conserved while the skeletal structure of the girdle that attaches those forelimbs to the body varies widely. Bony fish have a scapula and a cleithrum, though its homology to the mammalian clavicle is debated (Shubin et al., 2015). Many amphibians have clavicles that are widely recognized as homologous to the mammalian condition, as well as a fused scapulocoracoid (Jenkins & Goslow, 1983). Reptiles, including birds, have scapulae that articulate with bony furculae that are fused at the inferior end (Nesbitt, Turner, Spaulding, Conrad, & Norell, 2009). Many mammals possess clavicles to some degree, although they are often vestigial and

essentially nonfunctional. The presence of a clavicle of some form across marsupials and placental mammals and other tetrapods suggests that the lack of a clavicle is a derived trait.

Because the clavicle acts as a strut to distribute forces from the arm to the axial skeleton, its presence and morphology reflect the biomechanical forces that act on the forelimb during different kinds of locomotion. A pair of clavicles in the shoulder girdle enables the upper limbs to move outside the parasagittal plane (Jenkins, 1974), in contrast to bovids and other cursorial animals who lack clavicles and are unable to move their forelimbs perpendicular to their spine. Within primates, brachiators such as orangutans and gibbons have absolutely and relatively long, slender clavicles that are laterally twisted along the long axis. This morphology is an adaptation to these primates' superiorly oriented scapulae and tensile forces that act on the clavicles as the animals hang from their forelimbs. Arboreal quadrupedal primates, such as cercopithecins and colobines, have short, j-shaped clavicles and a deep fossa on the inferior side of the acromial end. As for fossils, Melillo (2016) interpreted *Australopithecus afarensis* clavicular morphology as evidence of increased manipulatory function of the upper limb.

The clavicle of *Homo sapiens* is notoriously variable and may well have the most developmental plasticity of any bone in the human skeleton. It is the first to begin ossification at around 9 weeks *in utero* and is one of the last bones to fully fuse between 20 and 30 years in age (Black & Scheuer, 1996). Developmentally, the clavicle is unique in having two primary ossification centers with different developmental origins. The medial end develops endochondrally from embryonic mesoderm while the lateral end develops intramembranously from embryonic neural crest (Matsuoka et al., 2005; Ogata & Uthoff, 1990). The extensive duration of ontogenetic development acting on these two centers of ossification means that the accumulation of biomechanical forces acting upon the shoulder girdle from before birth well into adulthood can influence adult clavicle morphology. Given this long ontogeny of the clavicle, the evolution of its morphology may well provide us with a window into a long period of the genetic and epigenetic forces experienced by the forelimbs of hominid individuals over geological time in addition to the broader insight to locomotory behavior.

In 2005, the lateral end of a hominid clavicle was recovered by AVS and other members of the Conservation Olduvai Project from Olduvai (Oldupai) Gorge, Tanzania. Following the historical practice of sequentially numbering hominid fossils from Olduvai Gorge, this clavicle is designated OH 89 (Figure 1). This is the second hominid clavicle discovered at Olduvai Gorge. The first, OH 48 was originally associated with the ~1.83 million-year-old OH 8, then later given its own number as its association with OH 8 was unclear (Day & Scheuer, 1989). OH 89 (Figure 1) preserves the lateral half of a right hominid clavicle.

OH 89 was found on the surface near the FLK North locality, eroding from upper Bed 1, just below tuff 1F (Figure 2). Tuff 1F has most recently been $^{40}\text{Ar}/^{39}\text{Ar}$ dated to 1.803 million years ago (Ma; Deino, 2012; Deino et al., 2021). Therefore, the minimum age of OH 89 is determined to be ~1.8 Ma. Given this geological age, OH 89 represents an important point of comparison between the earlier eastern African *A. afarensis* clavicles at 3.5 Ma, the approximately 1.95 Ma clavicular specimens from *Australopithecus sediba*, the 1.6 Ma clavicle of *Homo erectus* (KNM-WT-15000), and the clavicles of the more recent representatives of *Homo* from the Middle East and Eurasia. This time period is especially important given the morphological changes that occurred between *Australopithecus* and the appearance of *Homo*. The OH 89 clavicle, although isolated and fragmentary, provides an important data point in understanding the evolution of the hominid shoulder. OH 89 is now housed at the Arusha National Natural History Museum in Arusha, Tanzania.

Preservation and Anatomical Description of OH 89

OH 89 (Figure 1) has a maximum length of 87.9 mm and a maximum anteroposterior width of 20.4 mm. The acromial articular surface is missing, and the clavicle is broken around midshaft. There is a distinct subclavian groove, and the clavicle is roughly crescent shaped in cross section near the break. The rugosity for the deltoid is distinct and relatively more robust than that of the trapezius. The conoid tubercle appears as more of a rugosity than a tubercle.




Figure 1 - OH 89 in superior view (top) and inferior view (bottom).




Figure 2 - Map showing the location of OH 89 at Olduvai Gorge and the stratigraphic position of the fossil (left), with a map of Africa showing the geographic location of Olduvai Gorge (right). The stratigraphy is based on Hay (1976), Deino (2012), Deino et al. (2021), Deiz-Martín et al. (2015), McHenry & Stanistreet (2018), Njau et al. (2021), and Stanistreet et al. (2020).

Materials

We compared OH 89 to a sample of extant human and non-human primate and fossil hominid clavicles. Our extant sample consists of adult individuals from *Gorilla gorilla* (n=30), *Pan troglodytes* (n=31), *Homo sapiens* (n=25), and *Papio hamadryas* (n=7). We only included specimens that had no evidence of trauma or pathology. All extant non-human individuals come from wild-shot animals that are part of the Hamann-Todd Osteological Collection housed at the Cleveland Museum of Natural History (CMNH).

All 25 modern human individuals are housed in the Hamann-Todd Collection at CMNH. These skeletal remains are from people who died in the Cleveland area in the late 19th and early 20th centuries and whose bodies were unclaimed from hospitals and morgues (Cobb, 1933). These people therefore disproportionately represent low-income groups and, given the standards of medical ethics at the time, most likely did not agree to donate their remains to science.

All of these skeletal collections present complicated ethical questions (e.g., Walker, 2007; de la Cova, 2019; Squires and García-Mancuso, 2021). As our aim for this research is to better understand human and non-human primate biology from the broad perspective of geological time, we intentionally do not consider the ethnic or personal identification information of anyone included in this study. For the people whose clavicles were measured as part of in our study, we deeply appreciate their inclusion as representatives of *Homo sapiens*. For the non-human primates included in this study, we hope that using their remains to better understand primate biology at least derives some good from the sacrifice of wild animals.

Fossil clavicles included in this study are shown in Table 1. All measurements from fossils were taken on research quality casts of the original fossils, housed in the Human Evolution Research Center at UC Berkeley and the Cleveland Museum of Natural History.

Table 1. Fossil clavicles included in analyses.

Specimen Number	Side	Species	Date	References
KNM-WT 15000	R	<i>Homo erectus</i>	1.53 Ma	Walker & Leakey, 1993
KNM-ER 1808	R	<i>Homo erectus</i>	~1.5 Ma	Walker, Zimmerman & Leakey, 1982
OH 48	L	Unidentified	1.84 Ma	Napier, 1965
Krapina 158	L	Neanderthal	~130 Ka	Rink et al., 1995
KSD-VP-1/1	L	<i>Australopithecus afarensis</i>	3.6 Ma	Haile-Selassie et al., 2015

Methods

Linear Measurements

A series of 9 linear measurements were taken using Mitutoyo digital calipers: Maximum width of acromial end, maximum anteroposterior shaft width, maximum anteroposterior dimension, length of acromial rugosity, superoinferior width at midshaft, anteroposterior width at midshaft maximum superoinferior height of acromial region, anteroposterior width at conoid tubercle, and superoinferior width at conoid tubercle. Additionally, maximum length measurements were taken using an osteometric board. All linear measurements were taken directly on original specimens (for extant samples) or research quality casts (for fossil samples).

Length Estimate

Bilateral asymmetry of anthropoid clavicles is well documented, and results in right and left clavicles of the same individual exhibiting different lengths (Parsons, 1916; Schultz, 1937; Ray, 1959; Mays, Steele, & Ford, 1999; Voisin, 2001; Auerbach and Raxter, 2008; Abdel Fatah et al., 2012). We calculated an estimated length for OH 89 based off of the right clavicle using an ordinary least squares regression on midshaft size (√minimum midshaft width x perpendicular measurement), following Melillo (2016). These estimates use *Gorilla*, *Pan* and modern human individuals as a reference sample.

Curvature Measurement

Many researchers have quantified clavicular curvature through a range of different approaches, all of which have pros and cons (Olivier, 1951; Squyers & DeLeon, 2006; Voisin, 2006; Melillo, 2016; Barros, 2014). Linear measurements can quantify magnitude of curvature, but such analyses entirely lose the taxonomic shape profiles. Linear analyses are also unable to distinguish between a long clavicle with shallow curves and a short

clavicle with deep curves. Alternatively, 3D geometric morphometrics can be especially difficult to apply reliably to the clavicle because there are virtually no landmarks that are consistent across taxa, and those that are present are highly variable even within one species. Semi-landmarks may be utilized, but the reproducibility and accuracy of landmark placement is low and therefore less than ideal. 2D geometric morphometrics has most accurately captured clavicular shape and curvature thus far, but still has not been able to quantify the morphological differences as well as can be done by eye.

The methodology used here to assess clavicular curvature is partially modeled after that of Melillo (2016) but uses a simpler geometric approach instead of 2D geometric morphometric analyses. Given the preservation of OH 89, we only assess curvature from superior view. All specimens were photographed in a standard superior position with the acromial end horizontal to the ground. A curvilinear mid-line was drawn for each specimen that represents the sinusoidal shape of the bone. A composite is presented in Figure 3 to give the reader a sense of the range of variation in sinusoidal curvature in each of the taxa included in this study.




Figure 3 - Visualization of variation in clavicular curvature in comparative samples. Each midline curve was scaled to the same size and overlaid with others of the same taxon to create a “curvature cloud”. This visualization allows us to see the differences in overall shape of clavicular curvature between taxa as well as the variation within a group. All photos are scaled to the same size to better facilitate comprehension of shape without relation to size. Red lines indicate a female individual and black lines indicate a male individual.

Measurements of curves were taken in Adobe Illustrator© using the following steps (illustrated in Figure 4):

1. Midline curves were drawn on standardized photos of clavicles following the methodology of Melillo (2016).

2. The two endpoints of each midline curve were fitted to a horizontal line. The sternal and acromial endpoints are defined as the terminal midline point on each end of the clavicle.
3. Two circles were overlain on each clavicle using the “ellipse” tool in Adobe Illustrator®, one on the sternal and one on the acromial curve, such that the circle touches both endpoints of each curve and the maximum height of the curve.
4. A measurement of the radius of the sternal and acromial circles was taken. Midpoints for each circle were geometrically determined in Adobe Illustrator®, and radius of each circle were automatically calculated by Adobe Illustrator® in pixels, then converted to millimeters.




Figure 4 - An illustration of the methodology used to study curvature. See text for details.

These two curves (radius of sternal curve and radius of acromial curve) were then used to calculate a ratio of radii. Not only does this method capture both the length and depth of each curve, but it also allows for easier visualization of phylogenetic differences in morphology (see Results). Data are available at <https://doi.org/10.6084/m9.figshare.22050185.v1> (Taylor et al, 2023).

Analyses

We ran a series of standard univariate and bivariate analyses that include univariate descriptive statistics, a bivariate plot, and a principal component analysis (PCA). Univariate descriptive statistics were estimated using the *tapply* function of the *IRanges* package (Lawrence et al., 2013). Plots were produced using *ggplot2* (v3.3.0, Wickham, 2016) in the R statistical environment v3.6.1 (R Core Team, 2019) and edited in Adobe Illustrator CC 2019 (Adobe, Inc.). PCA includes all 11 measurements taken (the 9 linear measurements listed in section 4.1 and the two curve radius measurements described in section 4.3) and was performed using the *plotly* package (Sievert, 2020).

We also include a relatively new method of analysis for morphological studies – a t-distributed stochastic neighbor embedding analysis (t-SNE). This analysis was run using the *rtsne* package in R with a perplexity of 35 and 200 steps. t-SNE is a technique of visualizing high-dimensional data in two dimensions, somewhat similar to a PCA but with the capability of collapsing multi-dimensional data into a two-dimensional visualization (Krijthe, 2015; van der Maaten & Hinton, 2008). R code used in analyses can be found at <https://doi.org/10.6084/m9.figshare.22054280.v1> (Taylor, 2023). The goal of including this analysis was to determine if its use provides any additional information about fossil morphology when compared to a PCA. While the two analyses provide a similar way of visualizing data, t-SNE analyses collapse all of the data into two dimensions, rather than picking out two individual variables from the data and discarding the rest of the variables.

Results

Univariate descriptive statistics are presented in Table 2. To visualize the variation within and between taxa, maximum length is plotted against midshaft size for all extant and fossil data in Figure 5.




Figure 5 - A length regression on midshaft size based on the right clavicles of modern humans, *Gorilla*, and *Pan* individuals. *Pan* individuals are in dark blue, modern humans are in light blue, and *Gorilla* are in gold. Triangles indicate female and squares are male. OH 89's length estimate based on midshaft size is indicated by the grey oval, with a mean of 155.7 mm.

Table 2 - Descriptive statistics of fossils and extant comparative samples. Ranges are displayed as minimum value to maximum value. SI = Superoinferior and AP = Anteroposterior. All values are in mm except the ratio of radii. Ratio of radii is defined as the radius of the sternal curve divided by the radius of acromial curve.

	Maximum Length	Maximum width of acromial	Minimum AP Shaft Width	Maximum AP Dimension	Superoinferior Width at Midshaft	Maximum Superoinferior Height of Acromial Region	Anteroposterior Width at Conoid Tubercle	Superoinferior Width at Conoid Tubercle	Anteroposterior Width at Midshaft	Radius of Internal Curve	Radius of External Curve
<i>G. gorilla</i> (n = 30)	Mean 153.2 range 121.4 - 184.7	34 22.5 - 46.5	12.7 8.9 - 18.1	66.6 53.8 - 79.0	11.1 8.8 - 15.4	11.2 7.3 - 17.8	21.3 15.8 - 27.6	11.3 8.4 - 15.1	13.7 9.4 - 17.7	41.9 0 - 173.3	145.02 81.5 - 283.6
<i>P. troglodytes</i> (n = 31)	Mean 127.5 range 113.1 - 146.6	29.3 23.0 - 34.9	10.5 9.2 - 11.7	54.8 28.2 - 60.9	8.3 6.4 - 9.8	11.8 9.2 - 14.3	17.3 12.7 - 22.1	8.6 6.9 - 10.7	12.2 9.5 - 14.5	120.5 71.5 - 283.8	39.7 26.2 - 66.3
<i>H. sapiens</i> (n = 25)	Mean 144.2 range 127.4 - 158.8	31.5 26.2 - 37.0	11.6 8.7 - 14.0	62.5 54.8 - 68.6	9.8 7.8 - 13.1	11.4 7.9 - 14.5	17.7 15.1 - 21.9	10.9 8.3 - 15.2	12.1 9.0 - 14.8	91.6 69.4 - 135.8	49.1 29.9 - 74.0
<i>P. hamadryas</i> (n = 7)	Mean 73.24 range -	20.1 13.4 - 26.3	6.3 4.6 - 8.8	33.2 24.6 - 43.9	5.7 4.6 - 7.4	6.9 5.2 - 8.8	8.8 5.5 - 14.2	5.2 4.3 - 6.7	6.7 4.5 - 8.6	2.6 0 - 15.6	47.5 30.7 - 60.3
OH 48	Mean 148 range -	16.2 -	12.3 -	15.3 -	10.3 -	11.3 -	20.7 -	11.8 -	15.2 -	108.2 -	36.2 -
OH 89	Mean 155.7 range 153.3 - 158.0*	20.04 -	14.8 -	19.8 -	10.5 -	11.1 -	15.8 -	9.5 -	9.4 -	54.6 -	38.9 -
KNM-ER 1808	Mean 142.8 range -	14.7 -	10.6 -	12.3 -	9.3 -	9 -	15.8 -	8 -	12.2 -	90.2 -	35.4 -
KNM-WT 15000	Mean 130.5 range -	22.4 -	10.6 -	54.5 -	8.9 -	9 -	14.1 -	8.8 -	10 -	87.2 -	44.3 -
Krapina 158	Mean 126.7 range -	28.5 -	15.6 -	17.3 -	8.8 -	9.8 -	18.4 -	9 -	11.3 -	151.1 -	51.5 -
KSD-VP- 1/1	Mean 157.2 range -	30.6 -	15.3 -	24.6 -	9.6 -	10 -	20.3 -	10.4 -	16.4 -	62.2 -	43 -

Our length regression on midshaft size analysis estimates that OH 89 had a maximum length of 155.6 ± 2.4 mm. The fitted regression model was: Maximum length = $38.1713 + 9.3131 * \text{Midshaft size}$ ($R^2 = 0.73$, $p < 0.001$). Based on our regression on midshaft size, we estimate that the total length of OH 89 falls toward the upper end of modern human male variation in both maximum length and midshaft size. Compared to the other fossil clavicles included in our study, OH 89 is the largest in all dimensions.

Our quantification of clavicular curvature using circles captures the variation observed in bivariate plots (Figure 6), and PCA and t-SNE (Figure 7).




Figure 6 - Bivariate plots showing the ratio of radii vs. maximum length (top) and the radius of the acromial curve vs. maximum length (bottom).

For bivariate plots, both the ratio of radii and the radius of the acromial curve are used. Considering the fragmentary preservation of OH 89, we provide our estimate based on reconstructed length of the sternal curve, but error is high due to preservation, so the acromial curve is presented as well. While there is some overlap between humans and *Pan*, and humans and *Gorilla*, the three groups are mostly distinct. The PCA (Figure 7, left panel) appears to capture size and curvature in a similar manner to the bivariate plot (Figure 6). This quantitatively demonstrates that size and curvature profiles are the two most distinguishing characters in clavicles, according with the pattern discernible by eye.

Turning to the fossil data, we note that the fragmentary preservation of OH 89 requires that we interpret results with caution. With that said, the PCA shows minor overlap between chimpanzee and human individuals, with the hominid fossils falling within the two groups (Figure 7, left panel). OH 89 falls on the edge of the range of human variation in acromial curvature, and right on the edge between human and chimpanzee variation in the ratio of radii (Figure 6). Krapina 158 differs most apparently from *Homo sapiens* in comparison to the other fossils. The t-SNE results demonstrate a close affinity between humans and chimpanzees (Figure 7, right panel). In t-SNE analyses, the distance between clusters is not necessarily as informative as the clusters themselves. Thus, the plotting of several fossils on the top left does not necessarily indicate that those individuals are more similar to chimpanzees than humans, but does suggest that those individuals are more similar to each other than to any of the other groups. In this case, Figure 7, right panel shows that KNM-WT 15000 is most similar to humans, Krapina 158 to *Pan*, and that KNM-ER 1808, KSD-VP-1/1, OH 48, and OH 89 are most similar to each other.




Figure 7 - Left: Principal components analysis (PCA) showing fossil and extant clavicles. Right: t-distributed stochastic neighbor embedding (t-SNE) showing fossil and extant clavicles.

Discussion

Analyses of complete hominid clavicles have provided justification for claims about shoulder height, throwing ability, manual dexterity, and climbing ability (e.g., Larson, 2007; Roach & Richmond, 2015). The bilateral asymmetry of clavicles offers particular insight to behaviors, past and present (e.g., Abdel Fatah et al., 2012; Auerbach & Raxter, 2008; Trinkaus, Churchill, & Ruff, 1994). Given this relatively high degree of epigenetic influences on clavicular morphology compared to other regions of the skeleton, a fragmentary clavicle is a frustrating fossil to discover in isolation. Such a fossil is difficult to interpret taxonomically, and out of skeletal context, difficult to utilize in paleobehavioral studies. However, there is still information to be learned about human evolution from these fossils.

Our analysis of the fragmentary clavicle OH 89 provides clear evidence that a hominid died in the region of now- Olduvai Gorge 1.8 million years ago. This individual's clavicle was at the larger end of the variation we observe in *Homo sapiens* and is the largest of the hominid clavicles included in our analysis. While clavicle

length is not a good proxy for body size, it is a good predictor of the breadth of the individual's shoulders (Melillo et al., 2019). Our results indicate that OH 89 derives from an individual with a shoulder breadth similar to a large male today. With larger human comparative sample sizes, OH 89 would likely fall within the range of large humans.

One of the most defining morphological characteristics of the clavicle is its curvature. While this curvature is distinctive to the human eye (making the differentiation of hominoid clavicles fairly straight-forward qualitatively), quantifying these shape differences has been difficult. Multiple studies have attempted at quantifying clavicle curvature (Olivier, 1951; Voisin, 2006; Daruwalla, Courtis, Fitzpatrick, Fitzpatrick, & Millett, 2010; Abdel Fatah et al., 2012; Bernat et al., 2014; Squyers & DeLeon, 2015; Melillo, 2016; Barros, 2014). Here, we introduced a new method designed to be more easily applied to fragmentary fossils. Our method uses two circles to quantitatively capture the relative sinusoidal curvature of the clavicle. We find that the estimated sinusoidal curvature of OH 89's clavicle, as well as that of OH 48, is human-like relative to its length.

But OH 89 and OH 48 are clearly not identical to humans. Our PCA shows OH 89 and OH 48 at the intersection between humans and chimpanzees, along with KNM-WT 15000. A reasonable interpretation is that these three fossils are all part of the *Homo erectus* morphology, that chimpanzee clavicular morphology is ancestral and human morphology is derived. However, this interpretation is unsupported when we consider the anatomy of KSD-VP-1/1 which falls in the center of human variation (similar to Melillo's observation, 2016), and that the Neanderthal morphology of Krapina 158 is like that of chimpanzees. Researchers have noted the unique morphology of Neanderthal clavicles previously, so this aspect of our results is not entirely surprising (Mersey, Brudvick, Black, & Defleur, 2013; Voisin, 2006; Trinkaus, Holliday, & Auerbach, 2014; Rosas et al., 2016). However, the human-like morphology of the oldest clavicle included here (KSD-VP-1/1, 3.6 Ma), and the somewhat offset morphology of the 1.5 – 2 Ma clavicles suggest that evolution of the hominid clavicle has not been a constant movement towards a modern human morphology from a more chimpanzee morphology. This result echoes the conclusions reached by investigations of *Ardipithecus* anatomy (Lovejoy et al., 2009; Lovejoy and McCollum, 2010; White et al., 2015), showing that chimpanzees are distinctly derived compared to the chimpanzee-human last common ancestor (LCA) and therefore not an appropriate stand-in for the chimpanzee-human LCA.

Our t-SNE analysis, which relies on machine-learning (Figure 7), converts high-dimensional data into pairwise similarities, and thereby provides useful data visualization (Arora, Hu, & Kothari, 2018). The results of t-SNE analyses are designed to provide visual insight into the patterns formed by a data set, not provide quantitative results about the similarities and differences between groups. This type of analysis is usually employed in other disciplines with larger data sets, but it does provide insight into fossil morphology as illustrated here. We chose to utilize this analysis in an effort to determine if the conclusions of a t-SNE would be similar to a PCA, or if this analysis could provide further insight that the PCA does not. We found that most of the hominid clavicles cluster with each other, suggesting that they are more similar to each other than they are to any other clavicle included in the analysis. The two notable outliers to the hominid cluster are Krapina 158 and KNM-WT 15000. The Krapina clavicle is also somewhat of an outlier compared to the other hominids in its curvature profile, with a relatively larger sternal curve compared to the other fossils (Figure 6). Krapina 158 is also the shortest of the fossil clavicles, followed closely by KNM-WT 15000. The placement of KNM-WT 15000 closest to modern human in the t-SNE may be driven by the small size of this individual, given its sub-adulthood.

There are caveats to t-SNE, not the least of which is that it is usually used for big data sets with dozens to hundreds of variables, and therefore may be inappropriate for these much smaller datasets. However, it is fascinating that t-SNE returns an output that differentiates between our samples better than previously published analytical methods, and that these results compare favorably with what is visually obvious: human, chimpanzee, and gorilla clavicles look different from each other, KNM-WT 15000 falls most closely with modern humans, and that Krapina 158 is distinct from humans and other fossil hominids. These results support the conclusion by Voisin (2008) that OH 48 and KNM-WT 15000 have somewhat more pronounced sternal curves than acromial curves, though the results here place those two individuals on the edge of human variation.

In summary, our analyses demonstrate that OH 89 is, for the most part, on the edge of the range of variation of human clavicular variation despite its antiquity. Although researchers occasionally assign fragmentary, isolated postcrania to species, such a taxonomic identification is not possible for OH 89. Based on other fossil evidence, we know that around ~1.8 Ma, there were at least two distinct species of hominids living around Olduvai Gorge (Blumenschine et al., 2012; Clarke, 2012; Domínguez-Rodrigo et al., 2015). While it is impossible to decisively say what species OH 89 and OH 48 belonged to, or if they are even conspecific, the discovery of OH 89 does reveal that hominid clavicles show notable similarities to modern humans at least as far back as 1.8 Ma. This finding suggests that the evolutionary and biomechanical forces acting on the arms and shoulders of 1.8-million-year-old hominids are not entirely dissimilar to those acting on modern humans today.

Conclusion

OH 89, a 1.8 Ma hominid clavicle, falls within the range of modern humans in absolute size and clavicular curvature. This finding indicates that there has been little morphological change in the hominid clavicle in the last ~2 million years. It also suggests that shoulder breadth (though not necessarily body size) may have been similar to modern humans as far back as 1.8 Ma.

Acknowledgements

This project would not have been possible without the support and assistance of: Dr. H. Mshinda, Director General, Commission for Science and Technology (COSTECH); Mr. J. Paresso and Mr. J. Temba, Tanzanian Department of Antiquities; Ms. V. Ufunguo, Ngorongoro Conservation Area Authority; Dr. A. Mabulla and Mr. J. Kihyo, current and past directors of the National Museum of Tanzania; Dr. P. Msemwa, Dr. A. Kweka, and Dr. A. Gidna, Museum and House of Culture, Dar es Salaam, Tanzania; Mrs. F. Mangalu, Director of the National Natural History Museum, Arusha; OVPP's Olduvai field crew; Elias Bura, Theresa Grieco, Maneno Hussein, Odee Lyimo, Dan Mainoya, Justin Steven, Frank Mataro, Harry Mlaki, Leken Olle Moita, Mateo Oltusi, Singira Oltusi, Whitney Reiner, Kerio Zebedayo, Nairosi Zebedayo, Richard Bakari, and Ikoyo Singoi (aka Bruce Lee). Many thanks to the museums and staff who facilitate our access to the comparative skeletal collections, including L. Jellema from CMNH. Additionally, thank you to Yohannes Haile-Selassie for providing access to KSD-VP 1/1.

Preprint version 6 of this article has been peer-reviewed and recommended by Peer Community In Paleontology (<https://doi.org/10.24072/pci.paleo.100218>; Garcia, 2023).

Funding

This material is based upon work supported by the National Science Foundation under Grant No. BCS 1025263, NSF DGE 1752814, and was supported in part the European Research Council Advanced Grant Project 101054659—Tied2Teeth.

Conflict of Interest Disclosure

The authors acknowledge that one author (LJH) is a recommender for PCI Paleo.

Data Availability

Data used in this study are available at <https://doi.org/10.6084/m9.figshare.22050185.v1> (Taylor et al, 2023). R code is available at <https://doi.org/10.6084/m9.figshare.22054280.v1> (Taylor, 2023).

References

- Abdel Fatah, E. E., Shirley, N. R., Mahfouz, M. R., & Auerbach, B. M. (2012). A three-dimensional analysis of bilateral directional asymmetry in the human clavicle. *American journal of physical anthropology*, 149(4), 547-559. <https://doi.org/10.1002/ajpa.22156>
- Alemseged, Z., Spoor, F., Kimbel, W. H., Bobe, R., Geraads, D., Reed, D., & Wynn, J. G. (2006). A juvenile early hominin skeleton from Dikika, Ethiopia. *Nature*, 443(7109), 296-301. <https://doi.org/10.1038/nature05047>
- Arora, S., Hu, W., & Kothari, P. K. (2018). An analysis of the t-sne algorithm for data visualization. *Proceedings of the 31st Conference On Learning Theory*, PMLR 75:1455-1462. <https://proceedings.mlr.press/v75/arora18a.html>
- Auerbach, B. M., & Raxter, M. H. (2008). Patterns of clavicular bilateral asymmetry in relation to the humerus: variation among humans. *Journal of human evolution*, 54(5), 663-674. <https://doi.org/10.1016/j.jhevol.2007.10.002>
- Barros, A. (2014). Ontogeny, phylogeny and functional morphology of the hominoid shoulder girdle. (Doctoral dissertation). University College London. <https://discovery.ucl.ac.uk/id/eprint/1447948/>
- Bernat, A., Huysmans, T., Van Glabbeek, F., Sijbers, J., Gielen, J., & Van Tongel, A. (2014). The anatomy of the clavicle: A Three-dimensional Cadaveric Study. *Clinical anatomy*, 27(5), 712-723. <https://doi.org/10.1002/ca.22288>
- Black, S., & Scheuer, L. (1996). Age changes in the clavicle: from the early neonatal period to skeletal maturity. *International Journal of Osteoarchaeology*, 6(5), 425-434. [https://doi.org/10.1002/\(SICI\)1099-1212\(199612\)6:5<425::AID-OA287>3.0.CO;2-U](https://doi.org/10.1002/(SICI)1099-1212(199612)6:5<425::AID-OA287>3.0.CO;2-U)
- Blumenschine, R.J., Stanistreet, I.G., Masao, F.T. 2012. Olduvai Gorge and the Olduvai Landscape Paleoanthropology Project. *J Hum Evol* 63(2):247-250. <https://doi.org/10.1016/j.jhevol.2012.05.005>
- Carretero, J. M., Arsuaga, J. L., & Lorenzo, C. (1997). Clavicles, scapulae and humeri from the Sima de los Huesos site (Sierra de Atapuerca, Spain). *Journal of human evolution*, 33(2-3), 357-408. <https://doi.org/10.1006/jhev.1997.0128>
- Carlson, K. J., Green, D. J., Jashashvili, T., Pickering, T. R., Heaton, J. L., Beaudet, A., ... & Clarke, R. J. (2021). The pectoral girdle of StW 573 ('Little Foot') and its implications for shoulder evolution in the Hominina. *Journal of Human Evolution*, 102983. <https://doi.org/10.1016/j.jhevol.2021.102983>
- Clarke, R. J. (2012). A *Homo habilis* maxilla and other newly-discovered hominid fossils from Olduvai Gorge, Tanzania. *Journal of Human Evolution*, 63(2), 418-428. <https://doi.org/10.1016/j.jhevol.2011.11.007>
- de la Cova, C., 2019. Marginalized bodies and the construction of the Robert J. Terry anatomical skeletal collection: A promised land lost. In *Bioarchaeology of marginalized people* (pp. 133-155). Academic Press. <https://doi.org/10.1016/b978-0-12-815224-9.00007-5>
- Daruwalla, Z. J., Courtis, P., Fitzpatrick, C., Fitzpatrick, D., & Mullett, H. (2010). Anatomic variation of the clavicle: A novel three-dimensional study. *Clinical Anatomy: The Official Journal of the American Association of Clinical Anatomists and the British Association of Clinical Anatomists*, 23(2), 199-209. <https://doi.org/10.1002/ca.20924>
- Day, M. H. (1969). Early *Homo sapiens* Remains from the Omo River Region of South-west Ethiopia: Omo Human Skeletal Remains. *Nature*, 222 (5199), 1135. <https://doi.org/10.1038/2221135a0>
- Day, M. H. and Scheuer, J. L. (1989) Olduvai hominid 48: a clavicle. *Human Biology Budapest*, 19: 9– 14.
- Deino, A. L., Heil Jr, C., King, J., McHenry, L. J., Stanistreet, I. G., Stollhofen, H., ... & Toth, N. (2021). Chronostratigraphy and age modeling of Pleistocene drill cores from the Olduvai basin, Tanzania (Olduvai Gorge coring project). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 571, 109990. <https://doi.org/10.1016/j.palaeo.2020.109990>
- Deino, A. L. (2012). ⁴⁰Ar/³⁹Ar dating of Bed I, Olduvai Gorge, Tanzania, and the chronology of early Pleistocene climate change. *Journal of Human Evolution*, 63(2), 251-273. <https://doi.org/10.1016/j.jhevol.2012.05.004>
- Díez-Martín, F., Sánchez Yustos, P., Uribelarrea, D., Baquedano, E., Mark, D. F., Mabulla, A., ... & Domínguez-Rodrigo, M. (2015). The origin of the Acheulean: The 1.7 million-year-old site of FLK West, Olduvai Gorge (Tanzania). *Scientific reports*, 5(1), 1-9. <https://doi.org/10.1038/srep17839>

- Dirks, P. H., Kibii, J. M., Kuhn, B. F., Steininger, C., Churchill, S. E., Kramers, J. D., ... & Berger, L. R. (2010). Geological setting and age of *Australopithecus sediba* from southern Africa. *Science*, 328(5975), 205-208. <https://doi.org/10.1126/science.1184950>
- Domínguez-Rodrigo, M., Pickering, T.R., Almécija S., Heaton, J.L., Baquedano, E., Mabulla, A., Uribelarrea, D. 2015. Earliest modern human-like hand bone from a new >1.84-million-year-old site at Olduvai in Tanzania. *Nat Commun* 18(6):7987. <https://doi.org/10.1038/ncomms8987>
- Drapeau, M. S. M., Ward, C. V., Kimbel, W. H., Johanson, D. C., & Rak, Y. (2005). Associated cranial and forelimb remains attributed to *Australopithecus afarensis* from Hadar, Ethiopia. *Journal of Human Evolution*, 48(6), 593-642. <https://doi.org/10.1016/j.jhevol.2005.02.005>
- Fernandez-Jalvo, Y., Carlos Diez, J., Caceres, I. Rosell, Jordi (1999). Human cannibalism in the Early Pleistocene of Europe (Gran Dolina, Sierra de Atapuerca, Burgos, Spain). *Journal of human evolution*, 37(3-4), 591-622. <https://doi.org/10.1006/jhev.1999.0324>
- Feuerriegel, E. M., Green, D. J., Walker, C. S., Schmid, P., Hawks, J., Berger, L. R., & Churchill, S. E. (2017). The upper limb of *Homo naledi*. *Journal of Human Evolution*, 104, 155-173. <https://doi.org/10.1016/j.jhevol.2016.09.013>
- Garcia, N. (2023) A new method for measuring clavicular curvature. *Peer Community in Paleontology*, 100218. <https://doi.org/10.24072/pci.paleo.100218>
- García-González, R., Rodríguez, L., Salazar-Fernández, A., Arsuaga, J. L., & Carretero, J. M. Updated study of adult and subadult pectoral girdle bones from Sima de los Huesos site (Sierra de Atapuerca, Burgos, Spain). *Anatomical and age estimation keys. The Anatomical Record*. <https://doi.org/10.1002/ar.25158>
- Gómez-Olivencia, A., Quam, R., Sala, N., Bardey, M., Ohman, J. C., & Balzeau, A. (2018). La Ferrassie 1: New perspectives on a “classic” Neandertal. *Journal of Human Evolution*, 117, 13-32. <https://doi.org/10.1016/j.jhevol.2017.12.004>
- Green, D. J. (2020). Scapula, clavicle, and proximal humerus. In: *Hominin Postcranial Remains from Sterkfontein, South Africa*, Zipfel, B; Richmond, B.G., and Ward, C.V (eds). 1936-1995, 37. <https://doi.org/10.1093/oso/9780197507667.003.0005>
- Haile-Selassie, Y., Latimer, B. M., Alene, M., Deino, A. L., Gibert, L., Melillo, S. M., ... & Lovejoy, C. O. (2010). An early *Australopithecus afarensis* postcranium from Woranso-Mille, Ethiopia. *Proceedings of the National Academy of Sciences*, 107(27), 12121-12126. <https://doi.org/10.1073/pnas.1004527107>
- Hay, R. L. (1975). Olduvai Gorge: stratigraphy, age and environments. In *Abstract of communication at Geological Society Symposium: Geological Background to Fossil Man*.
- Heim, J.L. (1982). Les hommes fossils de La Ferrassie. II. Les squelettes d'adultes: squelettes de membres. Masson, Paris. <https://gallica.bnf.fr/ark:/12148/bpt6k33271664>
- Jenkins Jr, F. A. (1974). The movement of the shoulder in clavicate and aclavicate mammals. *Journal of Morphology*, 144(1), 71-83. <https://doi.org/10.1002/jmor.1051440105>
- Jenkins Jr, F. A., & Goslow Jr, G. E. (1983). The functional anatomy of the shoulder of the savannah monitor lizard (*Varanus exanthematicus*). *Journal of Morphology*, 175(2), 195-216. <https://doi.org/10.1002/jmor.1051750207>
- Krijthe JH (2015). Rtsne: T-Distributed Stochastic Neighbor Embedding using Barnes-Hut Implementation. R package version 0.16, <https://github.com/jkrijthe/Rtsne>.
- Larson, S. G. (2007). Evolutionary transformation of the hominin shoulder. *Evolutionary Anthropology: Issues, News, and Reviews: Issues, News, and Reviews*, 16(5), 172-187. <https://doi.org/10.1002/evan.20149>
- Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, Morgan M, Carey V (2013). “Software for Computing and Annotating Genomic Ranges.” *PLoS Computational Biology*, 9. <https://doi.org/10.1371/journal.pcbi.1003118>
- Lordkipanidze, D., Jashashvili, T., Vekua, A., De León, M. S. P., Zollikofer, C. P., Rightmire, G. P., ... & Rook, L. (2007). Postcranial evidence from early Homo from Dmanisi, Georgia. *Nature*, 449(7160), 305-310. <https://doi.org/10.1038/nature06134>

- Lovejoy, C. O., Johanson, D. C., & Coppens, Y. (1982). Hominid upper limb bones recovered from the Hadar formation: 1974–1977 collections. *American Journal of Physical Anthropology*, 57(4), 637–649. <https://doi.org/10.1002/ajpa.1330570409>
- Lovejoy, C.O., McCollum, M.A. Spinopelvic pathways to bipedality: why no hominids ever relied on a bent-hip-bent-knee gait. *Philos Trans R Soc Lond B Biol Sci.* 2010 Oct 27;365(1556):3289–99. <https://doi.org/10.1098/rstb.2010.0112>.
- Lovejoy, C. O., Suwa, G., Simpson, S. W., Matternes, J. H., & White, T. D. (2009). The great divides: *Ardipithecus ramidus* reveals the postcrania of our last common ancestors with African apes. *Science*, 326(5949), 73–106. <https://doi.org/10.1126/science.1175833>
- van der Maaten, L., & Hinton, G. (2008). Visualizing data using t-SNE. *Journal of machine learning research*, 9(11). <https://jmlr.org/papers/v9/vandermaaten08a.html>
- Matsuoka, T., Ahlberg, P. E., Kessaris, N., Iannarelli, P., Dennehy, U., Richardson, W. D., ... & Koentges, G. (2005). Neural crest origins of the neck and shoulder. *Nature*, 436(7049), 347–355. <https://doi.org/10.1038/nature03837>
- Mays, S., Steele, J., & Ford, M. (1999). Directional asymmetry in the human clavicle. *International Journal of Osteoarchaeology*, 9(1), 18–28. [https://doi.org/10.1002/\(SICI\)1099-1212\(199901/02\)9:1<18::AID-OA455>3.0.CO;2-A](https://doi.org/10.1002/(SICI)1099-1212(199901/02)9:1<18::AID-OA455>3.0.CO;2-A)
- McHenry, L. J., & Stanistreet, I. G. (2018). Tephrochronology of Bed II, Olduvai Gorge, Tanzania, and placement of the Oldowan–Acheulean transition. *Journal of Human Evolution*, 120, 7–18. <https://doi.org/10.1016/j.jhevol.2017.12.006>
- Melillo, S. M. (2016). The shoulder girdle of KSD-VP-1/1. In *The postcranial anatomy of Australopithecus afarensis* (pp. 113–141). Springer, Dordrecht. https://doi.org/10.1007/978-94-017-7429-1_6
- Melillo, S., Gunz, P., Coqueugniot, H., Reske, S., & Hublin, J. J. (2019). Structural effects of variation in the human clavicle. *American Journal of Physical Anthropology*, 168(4), 687–704. <https://doi.org/10.1002/ajpa.23787>
- Mersey, B., Brudvik, K., Black, M. T., & Defleur, A. (2013). Neanderthal axial and appendicular remains from Moula-Guercy, Ardèche, France. *American journal of physical anthropology*, 152(4), 530–542. <https://doi.org/10.1002/ajpa.22388>
- Nesbitt, S. J., Turner, A. H., Spaulding, M., Conrad, J. L., & Norell, M. A. (2009). The theropod furcula. *Journal of Morphology*, 270(7), 856–879. <https://doi.org/10.1002/jmor.10724>
- Njau, J. K., Toth, N., Schick, K., Stanistreet, I. G., McHenry, L. J., & Stollhofen, H. (2021). The Olduvai Gorge Coring Project: Drilling high resolution palaeoclimatic and palaeoenvironmental archives to constrain hominin evolution. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 561, 110059. <https://doi.org/10.1016/j.palaeo.2020.110059>
- Ogata, S., & Uthoff, H. K. (1990). The early development and ossification of the human clavicle—an embryologic study. *Acta Orthopaedica Scandinavica*, 61(4), 330–334. <https://doi.org/10.3109/17453679008993529>
- Olivier, G. (1951). Anthropologie de la clavicule. *Bulletins et Mémoires de la Société d'Anthropologie de Paris*, 2(4), 121–157. <https://doi.org/10.3406/bmsap.1951.2889>
- Parsons, F. G. (1916). On the proportions and characteristics of the modern English clavicle. *Journal of Anatomy*, 51(Pt 1), 71.
- Pearson, O. M., Royer, D. F., Grine, F. E., & Fleagle, J. G. (2008). A description of the Omo I postcranial skeleton, including newly discovered fossils. *Journal of Human Evolution*, 55(3), 421–437. <https://doi.org/10.1016/j.jhevol.2008.05.018>
- R Core Team. (2019). R: A language and environment for statistical computing. Retrieved from <https://www.R-project.org/>
- Radović, J., & Wolpoff, M. H. (1988). The Krapina hominids: an illustrated catalog of skeletal collection. Zagreb: Mladost for Croatian Natural History Museum. 118 p.
- Ray, L. J. (1959). Metrical and non-metrical features of the clavicle of the Australian Aboriginal. *American Journal of Physical Anthropology*, 17(3), 217–226. <https://doi.org/10.1002/ajpa.1330170307>

- Roach, N. T., & Richmond, B. G. (2015). Clavicle length, throwing performance and the reconstruction of the *Homo erectus* shoulder. *Journal of human evolution*, 80, 107-113. <https://doi.org/10.1016/j.jhevol.2014.09.004>
- Rosas, A., Rodriguez-Perez, F. J., Bastir, M., Estalrich, A., Huguet, R., García-Taberner, A., ... & de la Rasilla, M. (2016). Adult Neandertal clavicles from the El Sidrón site (Asturias, Spain) in the context of *Homo* pectoral girdle evolution. *Journal of human evolution*, 95, 55-67. <https://doi.org/10.1016/j.jhevol.2016.03.005>
- Schultz, A. H. (1937). Proportions, variability and asymmetries of the long bones of the limbs and the clavicles in man and apes. *Human Biology*, 9(3), 281-328.
- Senut, B., Nakatsukasa, M., Kunitatsu, Y., Nakano, Y., Takano, T., Tsujikawa, H., ... & Ishida, H. (2004). Preliminary analysis of *Nacholapithecus* scapula and clavicle from Nachola, Kenya. *Primates*, 45(2), 97-104. <https://doi.org/10.1007/s10329-003-0073-5>
- Shubin, N., Daeschler, E. B., & Jenkins Jr, F. A. (2015). Origin of the tetrapod neck and shoulder. *Great transformations in vertebrate evolution*, 63-76. <https://doi.org/10.7208/chicago/9780226268392.003.0004>
- Sievert C (2020). Interactive Web-Based Data Visualization with R, plotly, and shiny. Chapman and Hall/CRC. <https://doi.org/10.1201/9780429447273>
- Squires, K. and García-Mancuso, R., 2021. Ethical challenges associated with the study and treatment of human remains in anthropological sciences in the 21st century. *Revista Argentina de Antropología Biológica*, 23(2). <https://doi.org/10.24215/18536387e034>
- Squyres, N., & DeLeon, V. B. (2015). Clavicular curvature and locomotion in anthropoid primates: A 3D geometric morphometric analysis. *American journal of physical anthropology*, 158(2), 257-268. <https://doi.org/10.1002/ajpa.22785>
- Stanistreet, I. G., Stollhofen, H., Deino, A. L., McHenry, L. J., Toth, N. P., Schick, K. D., & Njau, J. K. (2020). New Olduvai Basin stratigraphy and stratigraphic concepts revealed by OGCP cores into the Palaeolake Olduvai depocentre, Tanzania. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 554, 109751. <https://doi.org/10.1016/j.palaeo.2020.109751>
- Taylor, Catherine (2023). OH89_RCode. figshare. Online resource. <https://doi.org/10.6084/m9.figshare.22054280.v1>
- Taylor, Catherine; Masao, Fidelis; Njau, Jackson; Songita, Agustino Venance; Hlusko, Leslea (2023). OH89_Data.csv. figshare. Dataset. <https://doi.org/10.6084/m9.figshare.22050185.v1>
- Trinkaus, E. (1982). The Shanidar 3 Neandertal. *American Journal of Physical Anthropology*, 57(1), 37-60. <https://doi.org/10.1002/ajpa.1330570107>
- Trinkaus, E., Churchill, S. E., & Ruff, C. B. (1994). Postcranial robusticity in *Homo*. II: Humeral bilateral asymmetry and bone plasticity. *American journal of physical anthropology*, 93(1), 1-34. <https://doi.org/10.1002/ajpa.1330930102>
- Trinkaus, E. (2011). The postcranial dimensions of the La Chapelle-aux-saints 1 Neandertal. *American Journal of Physical Anthropology*, 145(3), 461-468. <https://doi.org/10.1002/ajpa.21528>
- Trinkaus, E., Holliday, T. W., & Auerbach, B. M. (2014). Neandertal clavicle length. *Proceedings of the National Academy of Sciences*, 111(12), 4438-4442. <https://doi.org/10.1073/pnas.1402439111>
- Vandermeersch, B. (1981). *Les Hommes Fossiles de Qafzel (Israel)*. Cahiers de Paleoanthropologie. Edition du C.N.R.S., Paris
- Voisin, J. L. (2001). Évolution de la morphologie claviculaire au sein du genre *Homo*. Conséquences architecturales et fonctionnelles sur la ceinture scapulaire. *L'anthropologie*, 105(4), 449-468. [https://doi.org/10.1016/S0003-5521\(01\)80048-0](https://doi.org/10.1016/S0003-5521(01)80048-0)
- Voisin, J. L. (2006). Clavicle, a neglected bone: morphology and relation to arm movements and shoulder architecture in primates. *The Anatomical Record Part A: Discoveries in Molecular, Cellular, and Evolutionary Biology: An Official Publication of the American Association of Anatomists*, 288(9), 944-953. <https://doi.org/10.1002/ar.a.20354>
- Voisin, J. L. (2008). The Omo I hominin clavicle: Archaic or modern?. *Journal of human evolution*, 55(3), 438-443. <https://doi.org/10.1016/j.jhevol.2008.06.001>

- Walker, A., Leakey, R., & Leakey, R. E. (Eds.). (1993). The nariokotome *Homo erectus* skeleton. Harvard University Press.
- Walker, P.L. (2007) 'Bioarchaeological Ethics: A Historical Perspective on the Value of Human Remains', in M.A. Katzenberg and S.R. Saunders (eds) *Biological Anthropology of the Human Skeleton*, pp. 3–40. New York: Wiley-Liss. <https://doi.org/10.1002/9780470245842.ch1>
- Walker, M. J., Ortega, J., Parmová, K., López, M. V., & Trinkaus, E. (2011). Morphology, body proportions, and postcranial hypertrophy of a female Neandertal from the Sima de las Palomas, southeastern Spain. *Proceedings of the National Academy of Sciences*, 108(25), 10087-10091. <https://doi.org/10.1073/pnas.1107318108>
- White, T.D., Lovejoy C.O., Asfar B., Suwa G. 2015. Neither chimpanzee nor human, *Ardipithecus* reveals the surprising ancestry of both. *Proceedings of the National Academy of Sciences* 112(16):4877-4884. <https://doi.org/10.1073/pnas.1403659111>
- Wickham, H. (2016). *ggplot2: elegant graphics for data analysis*. Springer. <https://doi.org/10.1007/978-0-387-98141-3>